## Supplementary material (1)

## Criteria for Rasmussen radiologic assessment.

Subjective	Points
A. Articular depression	
Not present	6
<5 mm	4
6–10 mm	2
>10 mm	0
B. Condylar widening	
Not present	6
<5 mm	4
6–10 mm	2
>10 mm	0
C. Angulation (valgus/varus)	
Not present	6
<10°	4
10-20°	2
>20°	0
Maximum	18
Excellent	18
Good	12-17
Fair	6-11
Poor	<6

Fragment Specific Fixation for Tibial Plateau Fractures with particular reference to arthroscopic assisted surgery.

## Supplementary material (2)

Table: (1): Tegner Lysholm knee score

Patient's

Every step (0)

name (or Clinician's name (or ref)

ref)

This questionnaire has been designed to give your therapist information as to how your knee pain has affected your ability to manage in everyday life. Please answer every question by placing a mark in the box that best describes your condition today.

## **During the past 4 weeks.....**

Section 1 –Limp
None (5)
Slight or periodical (3)
Severe and constant (0)
Section 3 – Pain
None (25)
Inconstant and slight during severe
exertion (20)
M 1 11 ' (15)
Marked during severe exertion (15)
Marked on or after walking more than
2 km (10)
Marked on or after walking less than 2 km (0)

Section 6 – Swelling	Section 5 –Locking
None (10)	No locking and no catching sensations (15)
On severe exertion (6)	Catching sensation but no locking (10)
On ordinary exertion (2)	LockingOccasionally (6)
Constant (0)	Frequently (2)
	Locked joint on examination (0)

Constant

Fragment Specific Fixation for Tibial Plateau Fractures with particular reference to arthroscopic assisted surgery.

Section 8 – Squatting	Section 7 - Stair-climbing
No problems (5)	No problems (10)
Slightly impaired (4)	Slightly impaired (6)
Not beyond 90° (1)	One step at a time (2)
Impossible (0)	Impossible (0)