

Research Article

Assessment of Depression and Anxiety Disorders in Hearing Impaired Children

Samira Z. Sayed*, Samir M. Moner*, Ahmed A. Mohamed**, and Amr M. Nabil*

* Department of Pediatrics, Faculty of Medicine, Minia University, Egypt

**Department of Phoniatics, Faculty of Medicine, Minia University, Egypt

Abstract

Hearing impairment is one of the commonest birth defects which may result in some psychological disorders in childhood. The objective of this study is to assess the psychopathological complications of hearing impaired children compared to normal hearing ones. This was a prospective study which was conducted on students of two schools (Al Amal School for the deaf and the dumb and Copts School) at Minia city, Minia governorate, Egypt. A total of forty children aged 7:12 years were classified into two groups, group (I) patients: Included 20 male children with different degrees of hearing impairment, group (II) control: Included 20 healthy male children with normal hearing. Psychometric evaluation, electroencephalography, audiological evaluation and psychological assessment were done for all children. The results showed that cases with EEG abnormalities, different degrees of anxiety were significantly higher in patients group compared to control ($p < 0.05$) however, IQ did not differ. The study revealed that hearing impairment is associated with EEG abnormalities and the presence of both anxiety and depression which are the most common psychological disorders in hearing impaired children.

Keywords: Depression, Anxiety, Psychological Disorders, Hearing Impaired, Children.

Introduction

Hearing Impairment (HI) is a broad term that refers to hearing losses of varying degrees, ranging from hard-of-hearing to total deafness, there are many classifications of hearing loss, one of these is classification according to type includes conductive, sensorineural, or mixed hearing loss⁽¹⁾. Hearing loss is defined in terms of decibels (dB) lost, as follows: Slight hearing loss (16-25 dB), mild hearing loss (26-40 dB), moderate hearing loss (41-70 dB), severe hearing loss (71-90 dB) and profound hearing loss (> 90 dB)⁽²⁾.

Hearing impairment is the commonest birth defect and has a significant impact on both individual and society, it is considered to be the most prevalent congenital defects in neonates and its prevalence is higher than other conditions found at birth⁽³⁾. Globally, it was estimated that over of 5% has different degrees of hearing loss, of these, 32 million children⁽⁴⁾.

In Egypt, the prevalence of HI in children was reported as 13.8%⁽⁵⁾ and 20.9%⁽⁶⁾. In addition, it has been reported that the prevalence of hearing impairment increases with increasing age⁽⁶⁾.

Hearing impairment lowers the quality of life and has some bad psychological, physical and social consequences⁽⁷⁾. Many studies suggest that hearing loss is associated with reduced cognitive functioning and incident dementia⁽⁸⁾. Also, hearing impaired children are known to have more mental health problems, problems at school, experience more depression, and demonstrate more difficulties with social interactions⁽⁹⁾. In addition, there is a significant association between hearing impairment and moderate to severe depression and it is also a strong cause of anxiety in children and adolescents⁽¹⁰⁾.

The objective of this study is to assess the psychopathological complications such as

anxiety and depression of hearing impaired children as compared to normal hearing ones.

Subjects and methods

Study design

This was a prospective study which was conducted on students of two schools (Al Amal School for the deafs and the dumbs and Copts School) at Minia city, Minia governorate, Egypt during the period from January to June 2016.

Forty children aged 7:12 years were included in this study, they were divided into two groups matched in age and sex as follow:

Group (I) Patients: Included 20 male children with different degrees of hearing impairment according to audiometry, this group was divided to two subgroups:

Group (Ia): Included 10 children with severe hearing impairment (Minimum heard sound at 71-90 dB).

Group (Ib): Included 10 children with profound hearing impairment (Minimum heard sound more than 90 dB).

Group (II) Control: Included 20 healthy male children with normal hearing.

Methods and instruments

All included children were subjected to:

a) Through history taking stressing on history of any infection, trauma or drugs, perinatal history to exclude congenital infection, full developmental history to collect important information (such as sitting up, crawling, walking and talking) and also, family history and consanguinity.

b) Clinical examination: Such as general condition, vital signs, psychological assessment and neurological examination to detect or exclude any other clinical diseases.

c) Psychometric evaluation: To measure (I.Q) of cases by an expert psychologist by using Stanford–Binet Intelligence Scale 5th Edition (Table, 1)⁽¹¹⁾.

Table (1): Stanford–Binet classification.

IQ Range ("deviation IQ")	IQ Classification
145–160	Very gifted or highly advanced
130–144	Gifted or very advanced
120–129	Superior
110–119	High average
90–109	Average
80–89	Low average
70–79	Borderline impaired or delayed
55–69	Mildly impaired or delayed
40–54	Moderately impaired or delayed

d) Electroencephalography (EEG): To detect presence or absence of any defect in brain electrical activity.

e) Audiological evaluation: It was done detect the different degrees of hearing impairment or to prove normal hearing.

f) Psychological (Depression and Anxiety) assessment questionnaires (included 2 questionnaires):

1) Child Depression Inventory questionnaire (CDI): It is a psychological assessment that rates the severity of symptoms related to depression or dysthymic disorder in children

The questionnaire was well translated into Arabic by a psychiatric expert at Minia University Hospital.

2) The State-Trait Anxiety Inventory (STAI): It is an introspective psychological inventory consisting of 40 self-report items pertaining to anxiety affect and a commonly used measure of trait and state anxiety. Higher (STAI) scores means higher levels of anxiety. It's goal was to compile a set of items that could measure anxiety at both poles of the normal affect curve (state vs. trait).

Data analysis

Data was entered and statistically analyzed using SPSS program, version 20. Quantitative data was presented as mean \pm Standard deviation (SD), qualitative data was presented as frequency (%). Analysis was done by Chi-square, one-way ANOVA, t-test and Mann-Whitney U test when necessary. Probability value (P. value) of (< 0.05) was considered as a significant.

Results

Regarding EEG findings (Table, 2), there was a significant increase in EEG abnormalities in patients group compared to control, 3 cases (30%) in severe HI group and 6 cases (60%) in profound HI group versus 2 cases (10%) in control group, however there was no significant difference between severe and profound HI subgroups regarding EEG abnormalities. The same trend of EEG results was obtained in anxiety and depression, the degree of both anxiety and depression was significantly higher in HI patients (either severe and profound) compared to controls (Table, 2). The results

revealed that IQ classification was not significantly affected by EEG findings, anxiety and depression in both HI patients or controls. In the same line with our findings, cases with moderate anxiety and depression had significantly higher EEG abnormalities in patients group than control. There was no significant relation between incidence of depression or anxiety and EEG abnormalities in the same group (Table, 3).

Discussion

Hearing impairment is one of the most common birth defects significantly affect persons and their relation with the society especially, in children. Globally, it is estimated that about 32 million children have hearing impairment⁽⁴⁾, however, it was reported in studies conducted in Egypt that HI prevalence in children was 13.8%⁽⁵⁾ and 20.9%⁽⁶⁾. Generally, hearing impaired children are highly vulnerable to bad psychosocial development, they face communicational deficits difficulty their social and emotional life⁽⁹⁾.

Table (2): EEG, anxiety and depression among groups.

Variable	Group (I) Patients		Group (II) Control (n=20)	P. value (Sig.)
	Group (Ia) Severe HI (n=10)	Group (Ib) profound HI (n=10)		
EEG				
Normal	7 (70%) ^b	4 (40%) ^b	18 (90%) ^a	0.014*
Abnormal	3 (30%)	6 (60%)	2 (10%)	
Anxiety				
Mild	4 (40%) ^b	3 (30%) ^b	15 (75%) ^a	0.011*
Moderate	6 (60%)	7 (70%)	5 (25%)	
Depression				
Mild	7 (70%) ^b	3 (30%) ^b	17 (85%) ^a	0.018*
Moderate	3 (30%)	7 (70%)	3 (15%)	

* Significant ($p < 0.05$).

a,b Values with different superscripts are significantly different.

Table (3): Relation between anxiety, depression and EEG findings in different groups.

Variable	Group (I) Patients		Group (II) Control		P. value (Sig.)
	Normal EEG (n= 11)	Abnormal EEG (n= 9)	Normal EEG (n=18)	Abnormal EEG (n=2)	
Anxiety					
Mild	5 (45.5%) ^b	2 (22.2%) ^b	13 (72.2%) ^a	2 (100%) ^b	0.011*
Moderate	6 (54.5%)	7 (77.8%)	5 (27.8%)	0	
Depression					
Mild	5 (45.5%) ^b	5 (55.6%) ^b	15 (83.3%) ^a	2 (100%) ^b	0.018*
Moderate	6 (54.5%)	4 (44.4%)	3 (16.7%)	0	

* Significant ($p < 0.05$).

a,b Values with different superscripts are significantly different.

Regarding the results of IQ, there was no obvious difference in IQ level between HI and normal children. Similar results were obtained by⁽¹⁴⁾. Many studies reported that children with HI had lower IQ level and also communication skills compared to normal ones⁽¹⁰⁾. Children IQ level was strongly correlated with mental health status which was affected to high extent with hearing impairment as it was reported with many authors^(13,15).

The present results indicated that EEG abnormalities were significantly higher in patients group compared to control group (P-value: 0.013). Similar findings were found by Bailly et al.,⁽¹⁸⁾ and Van et al.,⁽¹³⁾ who found that EEG abnormalities was higher in hearing impaired children than normal hearing ones. The reason is unclear but, it may be attributed to some adverse changes occurred in the brain function of hearing impaired children. In line with our findings, in a study by Remine and Brown;⁽¹⁹⁾ to evaluate hearing loss impacts on brain function, they reported "by EEG findings" that brain function was delayed by hearing loss. Also, they added that when hearing loss occurs, brain areas which are responsible for other senses (vision or touch) will actually take over the areas of the brain which normally process hearing. On the other hand, Dammeyer,⁽²⁰⁾ reported that no differences were found between hearing impaired children and normal children in EEG results.

Our results revealed that patients group had significant increase in the presence of anxiety and depression compared to control group (P-

value: 0.011) and (P-value: 0.018) for anxiety and depression respectively, however, no significant differences were found between severe HL and profound HL subgroups. Our results are in agreement with Kvam et al.,⁽²¹⁾ who found that levels of depression and anxiety were higher in hearing impaired children as compared to control ones. Also, Theunissen et al.,⁽¹⁰⁾ found similar results. The possible explanation of this result that the HI children had communication problems which make social and emotional contact be difficult and make them always worried about the others. In addition, Rostami et al.,⁽²²⁾ who investigated retrospectively the presence of anxiety and depression among the hearing impaired persons based on researches results, they found that hearing impaired people had high degrees of anxiety and insomnia. However, Theunissen et al.,⁽²³⁾ did not found significant difference between hearing impaired children and normal hearing ones as regard anxiety disorders. Also, it was reported that hearing impaired children have more depressive symptoms than normally hearing children⁽²⁴⁾.

Accordingly, Tambs,⁽²⁵⁾ found "in about 50,000 subjects" that hearing loss was associated with substantially reduced mental health ratings (greater degree of depression and self-esteem) among young persons. Also, some studies indicate that the prevalence of mental health problems is 2-3 times higher than among their hearing peers, ranging from 20 to 50%^(19,20). It is thought that the higher incidence of depression in hearing impaired children may be due to the communicative barriers within a hearing world,

different causes of hearing loss and or may be related to stigma and discrimination⁽²¹⁾.

The present results showed a significant relation between EEG abnormalities and the prevalence of depression (P-value: 0.018) and anxiety (P-value: 0.011) in HI children. These result were supported by those of Hurt et al.,⁽²⁶⁾ and Simkin et al.,⁽²⁷⁾ who found that there are some changes in EEG were related to depression and anxiety because of nausea or abdominal distress, derealization or depersonalization and paresthesias that related to panic attacks, altered connectivity and activity in frontal and anterior cingulate cortex networks⁽²⁸⁾ or altered inflammatory cytokine and growth factor levels⁽²⁹⁾. Also, it was reported that EEG alpha activity elevated during rest in depressed patients⁽³⁰⁾. Furthermore, Begi et al.,

Conclusion

In conclusion, hearing impairment is significantly associated with EEG abnormalities. Also, the presence of both anxiety and depression was significantly increased in hearing impaired children than normal healthy ones and these disorders (anxiety and depression) are the commonest psychological disorders in hearing impaired children. Also, depression and anxiety in children are preventable and curable. This study have some limitations, of these: small sample size, we could not include females. So, we recommend further studies with larger sample size to evaluate the psychopathological complications of hearing impaired children and to study the impact of other variables such as methods of education, parental factors and its effects the child.

Ethical considerations: The study protocol and all procedures performed were in accordance with the ethical standards of Faculty of medicine- Minia University, Egypt and the aim and nature of the study was explained for each parent before inclusion. An informed written consent was obtained from parents before enrollment.

Acknowledgements: The authors thank all participants who were included in this project and their relatives and thank all who facilitated the conduct of this study.

Conflict of interest: The authors declare that they have no conflict of interest.

Funding: None.

References

1. World Health Organization. International Classification of Functioning, Disability and Health: ICF. World Health Organization, 15-32; 2001.
2. Komazec Z, et al., "Comparison between auditory steady-state responses and pure-tone audiometry." *Vojnosanitetski pregled* 67.9: 761-765; 2010.
3. Smith R, Gooi A. Etiology of hearing impairment in children, 40-76; 2012.
4. World Health Organization: Deafness and hearing loss. <http://www.who.int/mediacentre/factsheets/fs300/en/>; 2014.
5. Hamid O, Shoukry T, and Shehata W. "Prevalence and pattern of hearing loss in children in house-held national survey in Egypt." *Ejentas* 11.2: 12-6; 2010.
6. Taha AA, Pratt SR, Farhat TM et al., Prevalence and risk factors of hearing impairment among primary-school children in Shebin El-kom District, Egypt. *Am J Audiol.* 19(1) 46-60; 2010.
7. Borton SA, Mauze E, Lieu JE. Quality of Life in Children with Unilateral Hearing Loss: A Pilot Study. *Am J Audiol.*; 19(1): 61-72; 2010.
8. Mudar RA, Husain FT. Neural Alterations in Acquired Age-Related Hearing Loss. *Front. Psychol.* 7:828; 2016.
9. Theunissen SC, Rieffe C, Kouwenberg M, Soede W, Briaire JJ, Frijns JHM. Depression in hearing-impaired children. *Int J Pediatr Otorhinolaryngol* 75(10): 1313-1317; 2011.
10. Theunissen SC, Rieffe C, Kouwenberg M, De Raeve LJ, Soede W, Briaire JJ, Frijns JH. Behavioral problems in school-aged hearing-impaired children: the influence of sociodemographic, linguistic, and medical factors. *Eur Child Adolesc Psychiatry.* 23(4):187-96; 2014.
11. Kaufman A. *IQ Testing 101*. New York: Springer Publishing; 2009. p. 112. ISBN 978-0-8261-0629-2; 2009.
12. Rego MF, Ivone D, Rui N. Hearing impairment and nightmares: a theoretical insight. *Springerplus.* 4: 786; 2015.
13. Van DB, Lyndal C, Harvey D. "Sensitivity of cortical auditory evoked potential

- detection for hearing-impaired infants in response to short speech sounds." *Audiology Research* 2.1; 2012.
14. Theunissen SC, Carolien R, Wim S, Jeroen JB, Lizet K, et al., Symptoms of Psychopathology in Hearing-Impaired Children. *Ear & Hearing*;36:e190–e198; 2015.
 15. Van Gent T, Goedhart A, Hindley P, Treffers PD. Prevalence and correlates of psychopathology in a sample of deaf adolescents. *J Child Psychol Psychiatry* 48(9):950–958; 2007.
 16. Wingfield A, Peelle JE. The effects of hearing loss on neural processing and plasticity. *Front. Syst. Neurosci.* 9:35; 2015.
 17. Peelle JE, Troiani V, Grossman M, Wingfield A. Hearing loss in older adults affects neural systems supporting speech comprehension. *J. Neurosci.* 31 12638–12643; 2011.
 18. Bailly D, Dechoulydelenclave MB, Lauwerier L. [Hearing impairment and psychopathological disorders in children and adolescents. Review of the recent literature]. *Encephale.* 29(4P t 1):329-37; 2003.
 19. Remine MD, Brown PM. Comparison of the prevalence of mental health problems in deaf and hearing children and adolescents in Australia. *Aust N Z J Psychiatry* 44: 351-357; 2010.
 20. Dammeyer J. Psychosocial development in a Danish population of children with cochlear implants and deaf and hard-of-hearing children. *J Deaf Stud Deaf Educ* 15(1):50–58; 2010.
 21. Kvam M, Loeb M, Tambs K. Mental health in deaf adults: symptoms of anxiety and depression among hearing and deaf individuals, *Journal of Deaf Studies and Deaf Education* 12 :1–7; 2007.
 22. Rostami M, Bahman B, Vahid B. Depression and Deaf Adolescents: A review. *Iranian Rehabilitation Journal*, Vol. 12, No. 19; 2014.
 23. Theunissen SC, Rieffe C, Kouwenberg M, et al. Anxiety in children with hearing aids or cochlear implants compared to normally hearing controls. *Laryngoscope*, 122, 654–659; 2012.
 24. Valentijn SA, van Boxtel MP, van Hooren SA et al., Change in sensory functioning predicts change in cognitive functioning: results from a 6-year follow-up in the maastricht aging study. *J. Am. Geriatr. Soc.* 53 374–380; 2005.
 25. Tambs K. Moderate effects of hearing loss on mental health and subjective well-being: results from the Nord-Trøndelag Hearing Loss Study. *Psychosom Med.;* 66(5):776-82; 2004.
 26. Hurt E, Arnold LE, Lofthouse N. Quantitative EEG neurofeedback or the treatment of pediatric attention-deficit/hyperactivity disorder, autism spectrum disorders, learning disorders, and epilepsy. *Child Adolesc Psychiatr Clin N Am.* 23(3):465-86; 2014.
 27. Simkin DR, Thatcher RW, Lubar J. Quantitative EEG and neurofeedback in children and adolescents: anxiety disorders, depressive disorders, comorbid addiction and attention-deficit/hyper-activity disorder, and brain injury. *Child Adolesc Psychiatr Clin N Am.* (3):427-64; 2014.
 28. Chang Y, Xu J, Shi N, Pang X, Zhang B, Cai Z. Dysfunction of preattentive visual information processing among patients with major depressive disorder. *Biological Psychiatry*, 69, 742 – 747; 2011.
 29. Fox MD, Liu H, Pascual-Leone A. Identification of reproducible individualized targets for treatment of depression with TMS based on intrinsic connectivity. *NeuroImage*, 66C, 151 – 160; 2012.
 30. Schmidt HD, Shelton RC, Duman RS. Functional biomarkers of depression: Diagnosis, treatment, and pathophysiology. *Neuropsychopharmacology*, 36, 2375–2394; 2011.